

University of Dundee

Expanding the substrates for a bacterial hydrogenlyase reaction

Lamont, Ciaran M.; Kelly, Ciarán L.; Pinske, Constanze; Buchanan, Grant; Palmer, Tracy; Sargent, Frank

Published in:
Microbiology

DOI:
[10.1099/mic.0.000471](https://doi.org/10.1099/mic.0.000471)

Publication date:
2017

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Lamont, C. M., Kelly, C. L., Pinske, C., Buchanan, G., Palmer, T., & Sargent, F. (2017). Expanding the substrates for a bacterial hydrogenlyase reaction. *Microbiology*, 163(5), 649-653.
<https://doi.org/10.1099/mic.0.000471>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Expanding the substrates for a bacterial hydrogenlyase reaction

Ciaran M. Lamont,[†] Ciarán L. Kelly,[‡] Constanze Pinske,[§] Grant Buchanan, Tracy Palmer and Frank Sargent*

Abstract

Escherichia coli produces enzymes dedicated to hydrogen metabolism under anaerobic conditions. In particular, a formate hydrogenlyase (FHL) enzyme is responsible for the majority of hydrogen gas produced under fermentative conditions. FHL comprises a formate dehydrogenase (encoded by *fdhF*) linked directly to [NiFe]-hydrogenase-3 (Hyd-3), and formate is the only natural substrate known for proton reduction by this hydrogenase. In this work, the possibility of engineering an alternative electron donor for hydrogen production has been explored. Rational design and genetic engineering led to the construction of a fusion between *Thermotoga maritima* ferredoxin (Fd) and Hyd-3. The Fd-Hyd-3 fusion was found to evolve hydrogen when co-produced with *T. maritima* pyruvate::ferredoxin oxidoreductase (PFOR), which links pyruvate oxidation to the reduction of ferredoxin. Analysis of the key organic acids produced during fermentation suggested that the PFOR/Fd-Hyd-3 fusion system successfully diverted pyruvate onto a new pathway towards hydrogen production.

Under anaerobic fermentative conditions, *Escherichia coli* performs formate-dependent hydrogen production [1]. This is catalysed by the formate hydrogenlyase (FHL) complex [2–4], which is a membrane-bound enzyme comprising [NiFe]-hydrogenase-3 (Hyd-3) and a formate dehydrogenase component encoded by *fdhF* [5]. FdhF is loosely attached to Hyd-3 via the HycB protein, which itself contains four [4Fe-4S] clusters [2]. The *E. coli* Hyd-3 isoenzyme is unusual for a nickel-containing hydrogenase as it is apparently tuned towards proton reduction rather than H₂ oxidation [2]. However, this makes Hyd-3 an attractive candidate for engineering hydrogen production activity.

FHL subunits share sequence similarity with the membrane-bound hydrogenases (MBH) from, for example, *Pyrococcus furiosus* [6]. The electron donor for *P. furiosus* MBH is not a formate but a reduced ferredoxin [6, 7], probably generated by pyruvate::ferredoxin oxidoreductase (PFOR) [8]. PFOR is a cytoplasmic enzyme that oxidizes pyruvate to generate CO₂, acetyl-CoA, and reduced ferredoxin with a midpoint potential (E_m) estimated at –500 mV [9].

In this work, pyruvate was explored as an alternative non-natural substrate for H₂ production from *E. coli* Hyd-3. A rational design approach was taken to covalently attach the

ferredoxin from *Thermotoga maritima* to Hyd-3 via the HycB subunit. *T. maritima* Fd and PFOR plasmids were readily available [10]. To begin, strains were constructed where the natural electron donor enzyme for FHL, FdhF, was genetically removed (Table 1) using an available $\Delta fdhF$ allele [11]. *In vivo* hydrogen production assays involved measuring the accumulation of H₂ in the headspace (10 ml) of anaerobic cultures (5 ml) in Hungate tubes containing 0.8% (w/v) glucose. Following incubation at 37 °C, H₂ was quantified using gas chromatography (Shimadzu GC-2014) with N₂ as carrier (25 ml min^{–1}). The *fdhF* mutation resulted in a reduction in H₂-evolution activity of 1000 times compared to the original parent strain (Fig. 1a, b). This *fdhF* mutant phenotype was repeated in a strain carrying a chromosomal *hycE*^{His} allele (Table 1, Fig. 1b).

FHL subunits share similarity with the respiratory NADH dehydrogenase encoded by *nuoA-N* [3, 12, 13]. A $\Delta nuoA-N$ allele, marked with apramycin resistance from pIJ773 [14], had no effect on the ability of *E. coli* FGB300 or FTD300 (Table 1) to grow under fermentative conditions or the amount of H₂ produced (Fig. 1a). Next, $\Delta fdhF$ and $\Delta nuoA-N$ alleles were combined in a single strain (MG300dZ) and the double deletion was found to reduce the residual H₂

Received 19 January 2017; Accepted 10 April 2017

Author affiliation: School of Life Sciences, University of Dundee, Dundee DD1 5EH, Scotland, UK.

*Correspondence: Frank Sargent, f.sargent@dundee.ac.uk

Keywords: metabolic engineering; genetic engineering; fermentation; bio-hydrogen; hydrogenase; pyruvate::ferredoxin oxidoreductase.

Abbreviations: Fd, ferredoxin; FHL, formate hydrogenlyase; Hyd-3, [NiFe]-hydrogenase-3; MBH, membrane-bound hydrogenase; PFL, pyruvate formate-lyase; PFOR, pyruvate::ferredoxin oxidoreductase.

[†]Present address: Oxford BioMedica, Windrush Court, Transport Way, Oxford OX4 6LT, UK.

[‡]Present address: Department of Life Sciences, Imperial College London, South Kensington, London SW7 2AZ, UK.

[§]Present address: Institute of Biology/Microbiology, Martin Luther University Halle-Wittenberg, Kurt-Mothes-Str. 3, 06120 Halle (Saale), Germany. One supplementary figure is available with the online Supplementary Material.

Table 1. Strains and plasmids used or constructed in this study

Strain	Relevant genotype/description	Source
MC4100	<i>E. coli</i> K-12: F ⁻ , <i>araD139</i> , $\Delta(\arg F-lac)U169$, <i>ptsF25</i> , <i>deoC1</i> , <i>relA1</i> , <i>flbB5301</i> , <i>rspL150</i>	[22]
FTD300	As MC4100, $\Delta nuoA-N::Apra^R$	This work
MG1655	<i>E. coli</i> K-12: F ⁻ , λ^- , <i>ilvG</i> , <i>rfb-50</i> , <i>rph-1</i>	[23]
FGB300	As MG1655, $\Delta nuoA-N::Apra^R$	This work
MG16dZ	As MG1655, $\Delta fdhF$	This work
MG300dZ	As MG1655, $\Delta fdhF$, $\Delta nuoA-N::Apra^R$	This work
MG059e1	As MG1655, <i>hycE^{His}</i>	[2]
MGE1dZ	As MG1655, <i>hycE^{His}</i> , $\Delta fdhF$	[11]
FTF2013	As MGE1dZ, $\Delta nuoA-N::Apra^R$, $\Delta hycAB::\varphi fd-hycB$	This work
FTF2015	As MGE1dZ, $\Delta nuoA-N::Apra^R$, $\Delta hycAB::\varphi fd-hycB$, P _{T5} $\varphi fd-hycB$	This work
Plasmids		
pREP4	<i>lacI⁺</i> (Kan ^R)	Roche
pUNI-PROM	A pT7.5 derivative carrying 103 bp <i>E. coli</i> <i>tatA</i> promoter (Amp ^R)	[24]
pUNI-Tm-POR	As pUNI-PROM with <i>T. maritima</i> PFOR operon (Amp ^R)	[10]
pUNI-Tm-Fd-POR	As pUNI-PROM encoding <i>T. maritima</i> Fd and PFOR (Amp ^R)	[10]




Fig. 1. A fusion between ferredoxin and Hyd-3 produces hydrogen *in vivo* in the presence of pyruvate-ferredoxin oxidoreductase (PFOR). (a) The parental strains, MG1655 and MC4100, together with derivatives lacking the *nuo* operon encoding NADH dehydrogenase ($\Delta nuoA-N$) MG16dZ and FTD300, and the strain MG059e1 (as MG1655, *hycE^{His}*), were grown anaerobically in M9 medium supplemented with 0.8 % (w/v) glucose for 24 h after which the OD₆₀₀ was measured and the H₂ content in the headspace quantified by gas chromatography. Error bars represent SEM ($n=3$). (b) Strains carrying $\Delta fdhF$ deletions were analysed in an identical manner to those described in panel (a); however, the data are plotted separately as the values are 1000 times lower. (c) Strains FTF2013 ($\varphi fd-hycB$) and FTF2015 ($\varphi fd-hycB$ under control of the T5 promoter) were transformed with pUNI-PROM, pUNI-Tm-POR (encoding *T. maritima* PFOR) or pUNI-Tm-Fd-POR (encoding *T. maritima* PFOR and ferredoxin). The FTF2015 strain also carries pREP4 encoding LacI. Anaerobic M9 medium with 0.8 % (w/v) glucose, 0.2 % (w/v) casamino acids, plus 1 mM IPTG (final concentration) where indicated, was used. Cultures were incubated for 24 h at 37 °C. (d) Depiction of the complete PFOR/ φ Fd-Hyd-3 system activated in *E. coli*.

production further still (Fig. 1b). It is therefore possible that the very low levels of residual H_2 produced in the *fdhF* mutants results from reversed electron transport through Hyd-2 [15].

Next, a $\Delta hycA::\varphi fd-hycB$ allele was generated that encoded a fusion of *T. maritima* Fd to HycB via an HA epitope tag. Also, to upregulate expression of this fusion, the synthetic T5 promoter, *lac* operator and ribosome binding site from strain FZBup [11] was included to give a $\Delta hycAB::P_{T5}\varphi fd-hycB$ allele. Two strains, FTF2013 and FTF2015, were constructed (Table 1) and *in vivo* H_2 evolution activity quantified (Fig. 1c). The FTF2013 and FTF2015[pREP4] strains were transformed with pUNI-PROM (empty control vector), pUNI-Tm-POR (encoding *T. maritima* PFOR) or pUNI-Tm-Fd-POR (encoding *T. maritima* PFOR and Fd) then grown at 37 °C for 24 h in anaerobic Hungate tubes

containing 5 ml M9 medium supplemented with 0.8 % (w/v) glucose and 0.2 % (w/v) casamino acids. The FTF2013 strain produced H_2 at basal levels regardless of the presence of plasmids (Fig. 1c). This basal level was mirrored in the FTF2015[pREP4]/pUNI-PROM strain (Fig. 1c). However, when the PFOR plasmid was introduced into FTF2015 [pREP4] hydrogen, evolution increased to $>40 \text{ nmol } H_2 \text{ OD}^{-1} \text{ ml}^{-1}$ (Fig. 1c). Moreover, the vector encoding both PFOR and extra Fd induced H_2 production to a maximal level of $>60 \text{ nmol } H_2 \text{ OD}^{-1} \text{ ml}^{-1}$ in the presence of IPTG (Fig. 1c).

The levels of the most common organic acids produced during mixed-acid fermentation were investigated for strains producing active Fd-Hyd-3/PFOR (Fig. 2). Strains were grown for 24 h in 16 ml LB medium supplemented with 0.8 % (w/v) glucose. Culture supernatants were then passed




Fig. 2. The influence of the Fd-Hyd-3 fusion and PFOR on fermentation products. FTF2013 ($\varphi fd-hycB$) and FTF2015 ($\varphi fd-hycB$ under control of the T5 promoter) were each transformed with pUNI-PROM, pUNI-Tm-POR (encoding *T. maritima* PFOR) or pUNI-Tm-Fd-POR (encoding *T. maritima* PFOR and ferredoxin). FTF2015 also carries pREP4. Cultures were grown anaerobically in 16 ml LB plus 0.8 % (w/v) glucose and 1 mM IPTG (final), when required, at 37 °C for 24 h. The spent fermentation broth was analysed by HPLC by loading 5 μ l on an Aminex HPX-87H organic acid column at 0.5 ml min⁻¹ and 55 °C and monitoring absorbance at 210 nm. Organic acid standard curves were used all with R^2 values greater than 99.90 %. Peaks corresponding to the retention times of (a) formate, (b) lactate, (c) pyruvate, (d) acetate, and (e) succinate were quantified and data normalized to original OD₆₀₀. Error bars represent SEM ($n=3$). Note that succinate could not be confidently determined (ND) in samples containing IPTG. Lane 1, virgin LB medium + IPTG; lane 2, virgin LB medium + IPTG; lane 3, MC4100 positive control; lane 4, FTF2013 + pUNI-PROM; lane 5, FTF2013 + PFOR; lane 6, FTF2013 + PFOR + Fd; lane 7, FTF2015 + pUNI-PROM; lane 8, FTF2015 + PFOR; lane 9, FTF2015 + PFOR + Fd; lane 10, FTF2015 + pUNI-PROM + IPTG; lane 11, FTF2015 + PFOR + IPTG; and lane 12, FTF2015 + PFOR + Fd + IPTG.

through a 0.2 µm filter and analysed with an Aminex HPX-87H organic-acid column at 55 °C and 0.5 ml min⁻¹. Organic acids were detected by A_{210 nm} and compared to standard curves. Representative HPLC traces are shown in Fig. S1 (available in the online Supplementary Material). The starting concentration of glucose added to the rich medium was 44 mM D-glucose and, under the growth conditions chosen, the MC4100 FHL-positive strain produced 1.5 mM OD₆₀₀⁻¹ of formate (Fig. 2a) compared with 30.6 mM OD₆₀₀⁻¹ for the FTF2013/pUNI-PROM strain (inactive for FHL). Importantly, when the PFOR, Fd and Fd-Hyd-3 system is produced at its maximum level, the extracellular formate was observed to drop back to 5.7 mM OD₆₀₀⁻¹, which is indicative of pyruvate being directed away from the endogenous pyruvate formate lyase (PFL) enzyme to the fusion protein.

Extracellular lactate levels were found to be high in FTF2013 (Fig. 2b). This may mean that the higher formate levels (Fig. 2a) are inhibiting PFL leading to an accumulation of pyruvate and thus extra substrate for lactate dehydrogenase. Indeed, pyruvate can be detected in the growth medium (Fig. 2c) and its level does follow that of formate and lactate in the FTF2013 mutant strains (Fig. 2c). Although pyruvate would not normally be located outside the cell, *E. coli* is known to possess a pyruvate exporter to balance metabolite levels [16], and so any extracellular pyruvate levels may also correlate somewhat with that inside the cell cytoplasm. Importantly, in all cases, when the PFOR, Fd and Fd-Hyd-3 system is maximally produced (FTF2015/pUNI-Tm-Fd-POR + IPTG), the balance of pyruvate/lactate/formate returns to the low levels seen in the FHL-positive strain (Fig. 2c).

Normally, acetate levels are linked to that of acetyl CoA via phosphate acetyltransferase and acetate kinase. The observed increase in extracellular acetate (Fig. 2d) may mean a concomitant increase in cytoplasmic acetyl CoA, which is normally competed for by the AdhE-dependent ethanol production pathway. This is feasible and could be a consequence of the increased activity of the NADH-dependent lactate dehydrogenase already noted for these strains, which would reduce the requirement for AdhE to recycle NAD⁺ and allow acetyl CoA to be used for ATP and acetate production instead.

Together, these data demonstrate the successful repurposing of *E. coli* Hyd-3 to accept electrons from a new substrate: reduced ferredoxin linked to pyruvate :: ferredoxin oxidoreductase. Examples of native [NiFe]-hydrogenase :: ferredoxin interactions are not common; however, *Synechocystis* sp. PCC 6803 does contain such a system [17]. Physical tethering of a ferredoxin to an [FeFe]-hydrogenase, as opposed to a [NiFe]-hydrogenase, from *Chlamydomonas reinhardtii* showed that photosystem I could be coupled directly to H₂ production [18, 19]. Similarly, photosynthesis-linked ferredoxins have been fused to cytochromes P450 [20]. Functional fusion of a ferredoxin to [NiFe]-hydrogenases has resulted in some activity *in vitro* [21];

however, the φ Fd-Hyd-3 enzyme described in these experiments is one new example of a functional fusion that is active in the living cell.

Funding information

This work was funded by the Biotechnology and Biological Sciences Research Council EASTBIO Doctoral Training Partnership (BB/J01446X/1) and by responsive mode awards BB/I02008X/1 and BB/L008521/1.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

1. Sargent F. The model [NiFe]-hydrogenases of *Escherichia coli*. *Adv Microb Physiol* 2016;68:433–507.
2. McDowall JS, Murphy BJ, Haumann M, Palmer T, Armstrong FA et al. Bacterial formate hydrogenlyase complex. *Proc Natl Acad Sci USA* 2014;111:E3948–E3956.
3. Böhm R, Sauter M, Böck A. Nucleotide sequence and expression of an operon in *Escherichia coli* coding for formate hydrogenlyase components. *Mol Microbiol* 1990;4:231–243.
4. Sawers RG, Ballantine SP, Boxer DH. Differential expression of hydrogenase isoenzymes in *Escherichia coli* K-12: evidence for a third isoenzyme. *J Bacteriol* 1985;164:1324–1331.
5. Boyington JC, Gladyshev VN, Khangulov SV, Stadtman TC, Sun PD. Crystal structure of formate dehydrogenase H: catalysis involving Mo, molybdopterin, selenocysteine, and an Fe₄S₄ cluster. *Science* 1997;275:1305–1308.
6. McTernan PM, Chandrayan SK, Wu CH, Vaccaro BJ, Lancaster WA et al. Intact functional fourteen-subunit respiratory membrane-bound [NiFe]-hydrogenase complex of the hyperthermophilic archaeon *Pyrococcus furiosus*. *J Biol Chem* 2014;289:19364–19372.
7. Silva PJ, van den Ban EC, Wassink H, Haaker H, de Castro B et al. Enzymes of hydrogen metabolism in *Pyrococcus furiosus*. *Eur J Biochem* 2000;267:6541–6551.
8. Ma K, Hutchins A, Sung SJ, Adams MW. Pyruvate ferredoxin oxidoreductase from the hyperthermophilic archaeon, *Pyrococcus furiosus*, functions as a CoA-dependent pyruvate decarboxylase. *Proc Natl Acad Sci USA* 1997;94:9608–9613.
9. Thauer RK, Jungermann K, Decker K. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol Rev* 1977;41:100–180.
10. Kelly CL, Pinske C, Murphy BJ, Parkin A, Armstrong F et al. Integration of an [FeFe]-hydrogenase into the anaerobic metabolism of *Escherichia coli*. *Biotechnol Rep* 2015;8:94–104.
11. McDowall JS, Hjärsing MC, Palmer T, Sargent F. Dissection and engineering of the *Escherichia coli* formate hydrogenlyase complex. *FEBS Lett* 2015;589:3141–3147.
12. Efremov RG, Sazanov LA. The coupling mechanism of respiratory complex I: a structural and evolutionary perspective. *Biochim Biophys Acta* 2012;1817:1785–1795.
13. Marreiros BC, Batista AP, Duarte AM, Pereira MM. A missing link between complex I and group 4 membrane-bound [NiFe] hydrogenases. *Biochim Biophys Acta* 2013;1827:198–209.
14. Gust B, Challis GL, Fowler K, Kieser T, Chater KF. PCR-targeted *Streptomyces* gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proc Natl Acad Sci USA* 2003;100:1541–1546.
15. Pinske C, Jaroschinsky M, Linek S, Kelly CL, Sargent F et al. Physiology and bioenergetics of [NiFe]-hydrogenase 2-catalyzed H₂-consuming and H₂-producing reactions in *Escherichia coli*. *J Bacteriol* 2015;197:296–306.
16. Kreth J, Lengeler JW, Jahreis K. Characterization of pyruvate uptake in *Escherichia coli* K-12. *PLoS One* 2013;8:e67125.

17. Gutekunst K, Chen X, Schreiber K, Kaspar U, Makam S et al. The bidirectional NiFe-hydrogenase in *Synechocystis* sp. PCC 6803 is reduced by flavodoxin and ferredoxin and is essential under mixotrophic, nitrate-limiting conditions. *J Biol Chem* 2014;289:1930–1937.
18. Yacoby I, Pochekailov S, Toporik H, Ghirardi ML, King PW et al. Photosynthetic electron partitioning between [FeFe]-hydrogenase and ferredoxin: NADP⁺-oxidoreductase (FNR) enzymes *in vitro*. *Proc Natl Acad Sci USA* 2011;108:9396–9401.
19. Eilenberg H, Weiner I, Ben-Zvi O, Pundak C, Marmari A et al. The dual effect of a ferredoxin-hydrogenase fusion protein *in vivo*: successful divergence of the photosynthetic electron flux towards hydrogen production and elevated oxygen tolerance. *Biotechnol Biofuels* 2016;9:182.
20. Mellor SB, Nielsen AZ, Burow M, Motawia MS, Jakubauskas D et al. Fusion of ferredoxin and cytochrome P450 enables direct light-driven biosynthesis. *ACS Chem Biol* 2016;11:1862–1869.
21. Yonemoto IT, Smith HO, Weyman PD. Designed surface residue substitutions in [NiFe] hydrogenase that improve electron transfer characteristics. *Int J Mol Sci* 2015;16:2020–2033.
22. Casadaban MJ, Cohen SN. Lactose genes fused to exogenous promoters in one step using a Mu-lac bacteriophage: *in vivo* probe for transcriptional control sequences. *Proc Natl Acad Sci USA* 1979; 76:4530–4533.
23. Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V et al. The complete genome sequence of *Escherichia coli* K-12. *Science* 1997; 277:1453–1462.
24. Jack RL, Buchanan G, Dubini A, Hatzixanthis K, Palmer T et al. Coordinating assembly and export of complex bacterial proteins. *EMBO J* 2004;23:3962–3972.

Edited by: G. H. Thomas

Five reasons to publish your next article with a Microbiology Society journal

1. The Microbiology Society is a not-for-profit organization.
2. We offer fast and rigorous peer review – average time to first decision is 4–6 weeks.
3. Our journals have a global readership with subscriptions held in research institutions around the world.
4. 80% of our authors rate our submission process as 'excellent' or 'very good'.
5. Your article will be published on an interactive journal platform with advanced metrics.

Find out more and submit your article at microbiologyresearch.org.